



**ESCOLA UNIVERSITÁRIA VASCO DA GAMA**

**MESTRADO INTEGRADO EM MEDICINA VETERINÁRIA**

**RETROSPECTIVE STUDY ON MASTITIS PATHOGENS AND THEIR  
SUSCEPTIBILITY TO ANTIBIOTICS IN DAIRY FARMS OF  
SÃO MIGUEL (AZORES) – ASSOCIATION WITH HERDS' HOLDINGS  
AND MANAGEMENT PRACTICES**

Andreia Filipa Gomes Vieira

**Coimbra, Abril 2014**



**ESCOLA UNIVERSITÁRIA VASCO DA GAMA**

**MESTRADO INTEGRADO EM MEDICINA VETERINÁRIA**

**RETROSPECTIVE STUDY ON MASTITIS PATHOGENS AND THEIR  
SUSCEPTIBILITY TO ANTIBIOTICS IN DAIRY FARMS OF  
SÃO MIGUEL (AZORES) – ASSOCIATION WITH HERDS' HOLDINGS  
AND MANAGEMENT PRACTICES**

**Coimbra, Abril 2014**

**Andreia Filipa Gomes Vieira**

**Aluna do Mestrado Integrado em Medicina Veterinária**

**Orientador Interno**

**Prof. Dra. Sofia Duarte**

**Co-orientador**

**Dra. Elisabete Martins**



## **Agradecimentos**

Muito me apraz utilizar esta parte do meu trabalho para agradecer a todos aqueles que no decorrer do meu percurso me apoiaram, incentivaram e ajudaram, às quais qualquer agradecimento me soará necessariamente curto e insuficiente, pela importância individual de cada um no todo.

As minhas orientadoras, Dra. Sofia Duarte e Dra. Elisabete Martins, pela ajuda desde o primeiro minuto, pela dedicação, paciência, ensinamentos e simpatia. Ao professor Nuno por toda a ajuda prestada na análise estatística.

A toda a equipa da Associação de Jovens Agricultores Micaelenses com um especial agradecimento a equipa de veterinários com quem tive o prazer de trabalhar e por quem fui recebida com uma simpatia excecional, a Dra. Luísa Soares um obrigada por todo o apoio, ajuda, ensinamentos, disponibilidade e amizade, ao Dr. Tiago Viveiros por toda a disponibilidade, por todos os ensinamentos, pela amizade e por todas as gargalhadas, a Dra. Mónica e ao Dr. Luís um muito obrigada por toda a ajuda prestada. Ao Renato Sardinha, pela disponibilidade dos dados e pela ajuda nas questões técnicas. E a Sónia um muito obrigada por tudo. Um muito obrigado também a equipa da Caproper em especial a Dra. Rosa Rebelo e a Dona Paula pela disponibilidade e pela forma fantástica como fui recebida.

Aos meus amigos de Coimbra, que sabem quem são, obrigada por todos estes momentos maravilhosos ao longo destes anos, em especial ao Edgar e a Carolina, pelo apoio incondicional dado em todos os momentos.

Aos meus amigos de sempre, que desde o primeiro minuto me acompanharam nesta jornada um obrigada muito especial por todo o apoio e incentivo ao longo do meu caminho.

Por último, mas sem dúvida o mais importante, a minha família. Aos meus pais por me demonstrarem que o trabalho e a perseverança compensam e por me apoiarem neste sonho de menina. A minha irmã Ana pelo apoio ao longo destes anos e a minha irmã Alice por me mostrar que é bom sonhar.

Um obrigado a todos os que de alguma forma tornaram isto possível.

## Table of Contents

<b>Agradecimientos .....</b>	<b>V</b>
<b>LIST OF FIGURES.....</b>	<b>VII</b>
<b>LIST OF TABLES .....</b>	<b>VII</b>
<b>LIST OF ABBREVIATIONS.....</b>	<b>VII</b>
<b>TITLE PAGE .....</b>	<b>1</b>
<b>ABSTRACT .....</b>	<b>2</b>
<b>1. INTRODUCTION.....</b>	<b>3</b>
<b>2. MATERIALS AND METHODS .....</b>	<b>4</b>
<b>2.1. Characterization of the region and farms.....</b>	<b>4</b>
<b>2.2. Sample and data collection .....</b>	<b>5</b>
<b>2.3. Statistical analysis .....</b>	<b>6</b>
<b>3. RESULTS AND DISCUSSION .....</b>	<b>6</b>
<b>3.1. Mastitis pathogens .....</b>	<b>6</b>
<b>3.2. Antibiotic susceptibility patterns.....</b>	<b>10</b>
<b>3.3. Relationship between udder pathogens, herd's holding and management practices.....</b>	<b>13</b>
<b>ACKNOWLEDGMENTS.....</b>	<b>16</b>
<b>CONFLICT OF INTEREST STATEMENT .....</b>	<b>16</b>
<b>REFERENCES .....</b>	<b>17</b>
<b>ANNEX.....</b>	<b>21</b>
<b>Annex 1 .....</b>	<b>22</b>
<b>Annex 2 .....</b>	<b>35</b>

## LIST OF FIGURES

Figure 1. Isolated agents in analyzed samples

Figure 2. Percentage distribution of isolated environmental agents ( $n=689$ ) in the analyzed milk samples.

Figure 3. Percentage distribution of isolated contagious agents ( $n=137$ ).

## LIST OF TABLES

Table 1. Number of bacterial isolates susceptible to each tested AB.

## LIST OF ABBREVIATIONS

AJAM - Associação de Jovens Agricultores Micaelenses

Antibiotic – AB

IMM - Intramammary

CNS – Coagulase-negative *Staphylococci*

PFGE- Pulsed-field gel electrophoresis

# **Retrospective study on mastitis pathogens and their susceptibility to antibiotics in dairy farms of São Miguel (Azores) – association with herds' holding and management practices**

A. Vieira<sup>a,\*</sup>, S. Duarte<sup>a,b</sup>, E. Martins<sup>a</sup>

<sup>a</sup>*Department of Veterinary Medicine, Escola Universitária Vasco da Gama, Av. José R. Sousa Fernandes, 3020-210 Coimbra, Portugal*

<sup>b</sup>*Group of Health Surveillance - Center of Pharmaceutical Studies, Faculty of Pharmacy of the University of Coimbra - Polo III, 3000-548 Coimbra, Portugal*

\*Corresponding author: A. Vieira

E-mail: andreiavieira\_medvet@hotmail.com

Phone: +351917548343; Fax: +351239437627

## **ABSTRACT**

Bovine mastitis is the most prevalent disease in the production of dairy cattle worldwide.

The present study aimed to characterize the main etiological agents of mastitis and the corresponding susceptibility patterns in 700 milk samples submitted to the Azorean regional laboratory in 2013. A questionnaire was also performed to determine the relationship of the isolated agents with different types of management practices and features of the dairy farms that produced the analyzed milk samples.

Environmental agents were the ones more frequently isolated namely *Escherichia coli* and *Streptococcus uberis*. Contagious agents were less frequently isolated, of which *Staphylococcus aureus* was the most frequently isolated. Amoxicillin plus clavulanic acid was the antibiotic combination to which a larger number of agents was susceptible to.

Some management practices were significantly associated with the isolation of certain bacteria in mastitic milk samples, namely the extension of the dry periods and the use of internal teat sealant.

## **KEY WORDS**

Antibiotic susceptibility; Bovine; Management; Mastitis; Pathogens; Azores' herds.



## 1. INTRODUCTION

Bovine mastitis is the most prevalent disease found in the production of dairy cattle worldwide and accounts for several negative effects (Seegers *et al.*, 2003).

This disease is characterized by an inflammation of the mammary gland, caused mostly by bacteria but might also have fungi and algae as causative agents (Spanamberg *et al.*, 2009; Peixoto *et al.*, 2010). Udder pathogens have been divided in environmental and contagious organisms based in the transfer of organisms from the reservoirs to the teats between milkings or in the transmission from the carrier cow or quarter to the teats of non-infected cows/quarters during the milking process, respectively (Blowey & Edmondson, 2010)..

Milk yield and composition may be altered only in the short time in less severe mastitis cases. However, in more severe cases the changes can remain in the long-term. These effects may even occur in the following lactation (Seegers *et al.*, 2003).

In Portugal, the only available estimation of the costs of mastitis in dairies, refer to the Northern mainland region of Entre-Douro e Minho and are about 249€/mastitis/year. The reasons that mostly contribute to these losses are mortality and premature culling (Aires, 2010).

Due to occurrence of mastitis, there is an increase of the somatic cell count in the milk. A healthy udder and consequently a low level of somatic cells, avoids pecuniary penalization of the producers and, simultaneously, benefits the consumer and the dairy industry since both will have a product of greater quality (Blowey & Edmondson, 2010).

Currently, antibiotic (AB) therapy remains an important component of herd health programs to control intramammary (IMM) infections in dairy cows (Guérin-Faubleé *et al.*, 2002). The knowledge of the type of agent, as well as the susceptibility profile to several AB agents, can improve the success rates of the IMM infection treatment as well as contribute to the reduction of bacterial resistance (Silva *et al.*, 2004).

It is well established that the indiscriminate use of AB agents to treat mastitis or other type of infections may produce an increased level of resistance of many bacteria (Silva *et al.*, 2004). AB resistances have become an increasing problem in human and veterinary medicine, which can embody serious implications in terms of public health, since it may cause failures in treatments, death and prolonged illness, as well as increased costs associated with the treatment (Rangel & Marin, 2009).

The agricultural sector in the Azorean archipelago accounts for 10% of the region's economy, of which 54% refers to milk production. About 80% of the milk produced in Azores is exported and represents 27% of the total Portuguese dairy production (Ponte, 2009). In the year of 2013, in the Azorean archipelago, 536,074,200 liters of milk were produced, of which 65.1% were produced in the island of São Miguel (SREA, 2004).

The present study was drawn in view of the significance of the dairy farms to the local and national economy, and the absence of similar studies applied to this important milk-producing region. The main objectives were to characterize the main etiological agents identified in the laboratory submitted milk samples, as well as to determine their relationship with different types of management practices and the features of the dairy farms of the Azorean island of São Miguel. Furthermore, AB susceptibility patterns of major mastitis pathogens isolated were assessed, based on the laboratory records of one of the local dairy farmers association, throughout the previous year of 2013.

## **2. MATERIALS AND METHODS**

### **2.1. Characterization of the region and farms**

The farms from which the animals originated were located in the island of São Miguel, also known as the “green island”. This is the largest island of the Portuguese archipelago of Azores (composed by nine islands), situated in the Atlantic Ocean with their respective coordinates 37°47'N and 25°30'O. S. Miguel is located in the Oriental group of the archipelago and has an area of about 746,82km<sup>2</sup>. The climate in this region typically features a mild temperature with a high humidity, usually overcast with a regularly distributed rainfall throughout the year (Sousa, 2007).

Pastures occupy about 90% of the agricultural area in São Miguel. Terrain has steep slopes that sometimes hamper the access to farms and difficult agricultural work. The soils are very rich in potassium while very deficient in phosphorus, and, in a smaller scale, in calcium and magnesium. Regarding the micronutrients, studies reveal scarcity in selenium, zinc, copper, cobalt and iodine (Sousa, 2007).

The feeding regime of dairy cows is based primarily in pasture grazing. Currently, in São Miguel, most animals are not housed and are rather in pasture the whole year, which means that they are exposed to adverse weather conditions. Nevertheless, a small number of farms have started to build covered collective or individual areas with sand or straw. Mobile outdoor milking equipment is the most widely used. It is during milking that most farmers supply concentrate feeds.

The predominant breed on all herds is Holstein-Friesian although some producers are incorporating the Jersey breed in their farms, as they are animals that are better adapted to high and sloping pastures due to their low weight and height (Sousa, 2007).

## **2.2. Sample and data collection**

Data was collected from one of the two existing farmers' associations on the island of São Miguel, the *Associação de Jovens Agricultores Micaelenses* (AJAM). The AJAM gathers almost half of the dairy farms of the island, providing technical support and veterinary assistance. However, the laboratory of the AJAM serves other farms, beyond the ones registered in the association.

Laboratory data were collected retrospectively, in January 2014, and accounted for results of all samples submitted to the AJAM laboratory throughout the year 2013, by 133 different farms. Data records on the laboratory's database for each of the total 700 individual samples included individual animal identification, farm identification and farm location, as well as microorganism identification based on culture and their corresponding susceptibility for the following AB agents: marbofloxacin (5µg), danofloxacin (5µg), cefoperazone (75µg), cefquinome (30µg), kanamycin (30µg), amoxicillin plus clavulanic acid (25µg), cefazolin (30µg), spiramycin (100µg), lincomycin (75µg), gentamicin (120µg), ampicillin (10µg), trimethoprim sulfa (1,25µg), pirlimycin (2µg) and tylosin (30µg). An disk additionally employed comprised three AB agents, penethamate hydroiodide, dihydrostreptomycin sulphate and framycetin sulphate. Records were not obtained for resistance patterns.

In addition to the laboratory records, a questionnaire was held concerning the characteristics and management practices of  $n=105$  holdings. Only AJAM-associated farms were considered given that the questionnaire was performed during the scheduled veterinary support visits to the

farms between September 5<sup>th</sup> and December 11<sup>th</sup> of 2013. Previously informed producers consented for both collection and publication of data. The total number of cows of all enrolled herds was approximately 6,000 and the average farm size was about 170 animals, being the minimum herd size 23 animals and the maximum 320. The questionnaire was aimed at various aspects, in particular: feeding regime, existence of cowshed and cow maternity, number of milking cows, duration of dry periods and use of internal teat sealants and IMM AB at drying off. Particular aspects of milking hygiene were also abridged, such as the use of gloves during milking procedures, pre- and post-dipping with teat disinfectants, type of milking machine and criteria for milking cows with mastitis.

### **2.3. Statistical analysis**

Available data were submitted to several exploratory analyzes through the Excel<sup>TM</sup> and SAS<sup>TM</sup> (Statistical package) programs. Frequencies were determined for factors studied through the program Proc Freq SAS (SAS Institute, 2004).

The probability of occurrence of the various environmental and contagious agents was analyzed by logistic regression, with the Proc Logistic (SAS). The following factors were also evaluated: region, type of exploration, dry periods, teat sealant, teat dipping, criteria for milking cows with mastitis, type of milking machine, period of the year and number of animals.


Statistical significance was defined previously for test results with a p-value <0.01 and <0.05, for the different tests.

The calculation of other percentages was performed with Microsoft Excel 2010 software.

## **3. RESULTS AND DISCUSSION**

### **3.1. Mastitis pathogens**

Overall, a single agent was isolated in 65% (455/700) of the analyzed samples, whereas in 25% (176/700) two agents were isolated and in 1% (7/700) three agents. No growth was observed in about 9% (62/700) of the milk samples (Figure 1).




**Figure 1.** Isolated agents in analyzed samples, (1) – Samples with single agent isolates; (2) – Samples with 2 agents isolated; (3) – Samples with 3 agents isolated; (4) – Samples with no growth.

Of all agents isolated ( $n=826$ ) through culture, the majority corresponded to environmental agents (83%;  $n=689$ ) and, nearly 17% ( $n=137$ ) to contagious agents. The main reservoir of mastitis environmental agents is the environment of the herds and the exposure of healthy quarters to these agents can occur at any time during the life of a cow. On the other hand, for contagious mastitis, the first reservoir is the cow itself, and exposure of uninfected mammary quarters to contagious agents is restricted to the milking process (Zadocks *et al.*, 2001). As mentioned before, most cows in the herds of São Miguel are at pasture throughout the year, where they are exposed to environmental agents. An overcrowded pasture also contributes to a greater contamination by environmental agents (Blowey & Edmondson, 2010).

Considering the environmental agents only ( $n=689$ ; Figure 2), *Escherichia coli* (*E. coli*) corresponded to 30% (207/689) of isolates, whereas it corresponded to circa 25% (207/826) of the total positive samples. It should be noted, however, that the occurrence of this agent can be underestimated due to the false negatives resulting from the short duration of the infection, the fast immune response and its ensuing rapid elimination (Morn, 2009; Blowey & Edmondson, 2010). The bacteria *E. coli* is ubiquitous, being the most prevalent agent in farms, since it is present in feces and can contaminate different materials, such as bedding and water. This agent causes mastitis when teats are exposed to fecal contaminated material. Thus prevalence

of *E. coli* in farms increases with poor hygiene or high animal density. Rainfall and low frequency of pasture rotation can also influence the extent of exposure to *E. coli* (Morn, 2009; Rangel & Marin, 2009). These features are present in the herds of São Miguel, since most of the cows are at pasture and the climate presents regular rainfall throughout the year.

Coagulase-negative staphylococci (CNS) were isolated in nearly 14% (95/689) of the total isolated environmental agents, corresponding to 11.5% (95/826) of the total positive samples. CNS comprises 9 species and 16 subspecies. These agents are usually found in teat skin, teat end and teat canal of animals (Thorberg *et al.*, 2009; Blowey & Edmondson, 2010).



**Figure 2.** Percentage distribution of isolated environmental agents ( $n=689$ ) in the analyzed milk samples (CNS, Coagulase-negative *Staphylococci*).

*Streptococcus uberis* (*S. uberis*) was isolated in 13.2% (91/689) of the total environmental agents' isolates, corresponding to 11% (91/826) of the total positive samples. *S. uberis* is the most common causative agent of mastitis in modern dairy herds according to Zadocks *et al.* (2007), although only the 5<sup>th</sup> most frequently isolated agent in the farms of São Miguel. The capacity of this agent to survive in the environment may be enhanced by the presence of hyaluronic acid capsule (Lopez-Benavides *et al.*, 2007). Bedding with sand and straw is associated with high levels of *S. uberis*, as once contaminated with milk, urine or feces, will allow bacterial proliferation. However, it can also be found in the mouth, vulva, groin and axilla of cows. Furthermore, this agent can also affect cows in pasture, since it is present in feces. Nevertheless, its frequency of isolation is not comparable with the one of *E.coli* (Blowey & Edmondson, 2010). Zadocks (2007) refer that *S. uberis* can behave either as an environmental agent or as a contagious agent. Indeed, it has been observed in several farms that many

animals within a herd had the same or closely related strain of *S. uberis*. This could be the result of a predominance of a determined strain in the environment or otherwise a contagious transmission. Rato *et al.* (2013) reported heterogeneity in pulsed-field gel electrophoresis (PFGE) patterns between *S. uberis* isolates of different farms; as, a patterns were specific in the same farm, a direct transmission of *S. uberis* among animals within the same farm can thus be suggested.

The findings of the present study, referring to the herds of the São Miguel island, reinforce the need to prevent mammary infections namely by environmental agents like *E. coli*. Measures to prevent environmental mastitis include rotating pasture, avoiding overstocking, paying attention to hygiene around feeding areas, and creating maternities in order to avoid contamination at the time of parturition (Blowey & Edmondson, 2010).


Regarding classical contagious agents ( $n=137$ ), and as displayed in Figure 3, *Staphylococcus aureus* (*S. aureus*) corresponded to almost 60% (82/137) of the total isolated contagious agents and to nearly 10% (82/826) of the total positive samples. IMM infections by this agent are chronic and subclinical, with periods of clinical mastitis. When established, infection is extremely hard to eliminate. As the excretion of this agent is intermittent, a negative culture does not necessarily imply that the cow is free of *S. aureus*. Therefore, it is possible that the real occurrence of this agent is higher, taking into account the possibility of intermittent negative results during culture. These infections usually are spread from cow to cow at milking (Morn, 2009; Blowey & Edmondson, 2010).

*Streptococcus agalactiae* (*S. agalactiae*) was isolated in 22.6% (31/137) of the total isolated contagious agents, corresponding to 3.8% (31/826) of the total positive samples and is a major contagious pathogen. Despite an observed trend of decreasing occurrence, in the past few years, *S. agalactiae* remains a significant cause of mastitis in herds with poor management (Merl *et al.*, 2003). Blowey & Edmondson (2010) reported that the isolation of this agent is a good indicator of hygiene breaching during milking.

*Corynebacterium bovis* was isolated in 15% (21/137) of the total isolated contagious agents, corresponding to 2.5% (21/826) of the total positive samples. This agent can be isolated from the teat canal (Blowey & Edmondson, 2010). Comparing two techniques of milk collection, Bexiga *et al.* (2010) reported that milk samples collected with a cannula surpassing the teat

canal decreased the isolation of *Corynebacterium bovis* when compared with milk samples collected with the standard technique, probably because the first technique avoids contamination of the milk samples with teat canal flora.

*Mycoplasma* was the less frequently isolated agent. Considering the total isolated contagious agents, *Mycoplasma* was found in 2.2% (3/137), which corresponded to 0.4% (3/826) of the total positive samples. According to Pinho (2013), the prevalence of *Mycoplasma* in dairy herds of Northwestern Portugal was determined as not higher than 3%.



**Figure 3.** Percentage distribution of isolated contagious agents ( $n=137$ ).

### 3.2. Antibiotic susceptibility patterns

AB susceptibility was determined in 842 isolates as summed up in Table 1. Amoxicillin plus clavulanic acid (aminopenicillins) was the AB to which a larger number of agents was sensible to, specifically 15.9% (134/842). Considering *E. coli* isolates only, 28.6% (73/255) were susceptible to marbofloxacin, 14% (36/255) to danofloxacin (which represents a sensitivity of 42.6% of strains to the tested fluoroquinolones), and almost 12% (30/255) were susceptible to kanamycin (of the aminoglycoside group). Previously, Saini *et al.* (2012) reported the resistance of this agent to kanamycin, ampicillin, streptomycin and tetracycline. Although in the present study nearly 12% of the *E. coli* isolates were susceptible to kanamycin, it is not possible to assume that remaining isolates were resistant to this drug, as resistance patterns were not available. Rocha (2013) reported resistance of this bacteria to amoxicillin plus clavulanic acid in 30% of herds evaluated in mainland Portugal. In the present study, susceptibility of the agents to



this combination of AB was only 8.6%, but again patterns of resistance were not available to allow comparison with mainland.

In relation to *S. uberis* isolates, 19.6% (22/112) were susceptible to kanamycin, 18.8% (21/112) to amoxicillin plus clavulanic acid and approximately 11.6% (13/112) to spiramycin, a macrolide. Rocha (2013) reported resistance of *S. uberis* to gentamicin (71%) and cloxacillin (22.1%) in herds from the mainland of Portugal. In the routine antibiograms performed in the laboratory of AJAM, cloxacillin is not included. Moreover, sensibility of *S. uberis* to gentamicin in the São Miguel Island was only 1%. According to Blowey & Edmonson (2010), the low efficacy treatment of *S. uberis* and long refractory period in the udder, may be due to factors such as resistance to phagocytosis, maintenance inside the cells where many ABs cannot act, permanency in the mammary lymph node, kept as a reservoir of infection.

The results obtained for *S. aureus* showed that 24.6% (16/65) of isolates were susceptible to amoxicillin plus clavulanic acid, 20% (13/65) to cefoperazone (a third-generation cephalosporin) and 16.9% (11/65) to tylosin (macrolide). Rocha (2013) refer an increased resistance to the AB combination of amoxicillin plus clavulanic acid from 2004 (0.7%) to 2012 (26.6%), in dairy cattle from the mainland of Portugal. Once more, as only the susceptibility patterns were available for the herds of São Miguel, comparison of resistance patterns is not possible. Due to the production of enzymes and endotoxins by *S. aureus* strains, damage is caused in mammary cells which results in fibrosis and abscess formation. Some strains can also form biofilms in the udder. Consequently, IMM infections by *S. aureus* rarely resolve spontaneously and are difficult to treat with ABs (Morn, 2009; Melchior, 2009). Nevertheless, in a first-time infection the cure rates may be improved (Blowey & Edmondson, 2010). Through Polymerase Chain Reaction (PCR) it is possible to detect the presence of AB resistance genes or presence of genes that have been correlated with biofilm formation in *S. aureus*. The presence of these genes decreases the treatment success (Melchior *et al.*, 2009).

*S. agalactiae* has a good response to antibiotic therapy and almost all antibiotics are effective according to Blowey & Edmondson (2010). In this study, 21.6% (11/51) of *S. agalactiae* isolates were sensible to amoxicillin plus clavulanic acid and spiramycin. Previously, Rocha (2013) reported the lowest AB resistance proportions (0 to 11.4%) of *S. agalactiae* in comparison with the remaining tested bacterial isolates.

**Table 1.** Number of bacterial isolates susceptible to each tested AB.

	<i>E. coli</i> (n=255)	<i>S. Non-agalactiae</i> (n=121)	<i>S. uberis</i> (n=112)	<i>Enterococcus</i> (n=168)	<i>S. aureus</i> (n=65)	CNS (n=43)	<i>C. bovis</i> (n=1)	<i>S. agalactiae</i> (n=51)	<i>S. dysgalactiae</i> (n=10)	Others (n=15)	Total (n=842)
<b>Marbofloxacin</b>	73	5	0	6	0	2	0	0	0	2	<b>88</b>
<b>Danofloxacin</b>	36	4	0	1	0	3	0	0	0	0	<b>44</b>
<b>Cefoperazone</b>	28	12	5	29	13	11	1	3	0	4	<b>106</b>
<b>Cefquinome</b>	19	2	1	1	4	3	0	0	0	0	<b>30</b>
<b>Kanamycin</b>	30	11	22	0	3	1	0	5	1	1	<b>75</b>
<b>Amoxicillin + clavulanic acid</b>	22	20	21	26	16	9	0	11	3	9	<b>134</b>
<b>Cefazolin</b>	1	3	12	0	2	0	0	2	3	3	<b>23</b>
<b>Spiramycin</b>	7	11	13	21	7	3	0	11	1	2	<b>76</b>
<b>Lincomycin</b>	18	24	9	50	4	3	0	10	1	3	<b>121</b>
<b>Gentamicin</b>	0	0	1	0	1	0	0	0	0	0	<b>2</b>
<b>Ampicillin</b>	4	10	10	7	1	4	0	2	0	0	<b>38</b>
<b>Trimethoprim</b>	6	5	3	8	1	2	0	2	1	1	<b>28</b>
<b>Disk with 3 AB *</b>	11	12	13	19	2	2	0	4	0	0	<b>63</b>
<b>Pirlimycin</b>	0	0	1	0	0	0	0	0	0	0	<b>1</b>
<b>Tylosin</b>	0	0	1	0	11	0	0	1	0	0	<b>13</b>

(AB, Antibiotic; CNS, Coagulase-negative *Staphylococci*; *S. Non-agalactiae*, *Streptococcus non-agalactiae*; *C. bovis*, *Corynebacterium bovis*; *S. dysgalactiae*, *Streptococcus dysgalactiae*; \*, comprising penethamate hydroiodide, dihydrostreptomycin sulphate and framycetin sulphate;)

The purpose of the treatment of mastitis is to guarantee that effective AB concentrations are reached at the site of infection. The most frequently used route of administration for treatment of mastitis is IMM, as a large concentration of ABs in the mammary gland is reached because the AB is administered directly into the site of infection. With parenteral administration, it is difficult to achieve and maintain a concentration in milk or tissue. Systemic treatment is recommended in some situations, like in clinical mastitis due to *S. aureus* (in combination with IMM treatment) and severe coliform mastitis (Pyorola, 2006). Some classes of ABs are not available in preparation for IMM administration in the Portuguese market, as is the case of quinolones and macrolides. In the present study, the overall AB susceptibility of isolated bacteria to quinolones was nearly 16%.

### **3.3. Relationship between udder pathogens, herd's holding and management practices**

According to results of the questionnaire carried out in 105 holdings, some management procedures were significantly associated with the isolation of certain bacteria in mastitic milk samples.

The use of internal teat sealants had a positive association with the milk samples tested positive for *E.coli* ( $p<0.05$ ). According to such result, holdings that always use internal teat sealants have a higher probability to have *E. coli* mastitis, whereas, holdings that rarely use internal teat sealants have less probability to have this type of mastitis. This finding is in contrast with previous studies that reported significantly fewer IMM infection using teat sealants (Berry & Hillerton, 2002). This result can also be explained by the fact that teat sealant is being mostly used by farms that have greater mastitis problems. In addition, if aseptic techniques are not warranted during sealant installation, pathogens present around the teat, in the environment or in the disinfectants and utensils can reach the teat ends and the teat canal, promoting infection (Huscley *et al.*, 2002). As such, in future studies it is necessary to ascertain in more detail this association.

Additionally, the herd size had a statistically significant association with isolation of *E. coli* and *Enterococcus* ( $p<0.01$ ). Regarding *E. coli*, the frequency of isolation increases as the number of animals in the herd increases, whereas for *Enterococcus*, the positivity of isolation decreases when the number of animals in the herd increases. Likewise, already previously, Rangel and

Marin (2009) reported that *E. coli* increases in farms featuring a greater animal density and larger herds might be in greater risk for pasture overcrowding.

Milking order of mastitic cows also showed a statistically significant association with *Enterococcus* ( $p < 0.05$ ) isolation. Statistical analysis showed that farms where cows with mastitis were left to be milked last were less prone to test positive for *Enterococcus*. This is in accordance with the recommended measures for prevention and control of mastitis in dairy cows (as recently reviewed by Blowey and Edmondson, 2010). But given that *Enterococcus* is not a contagious agent it is surprising that this association has not occurred with contagious agents, as infections with contagious agents usually are spread from cow to cow at milking.

Possible effects of dry period length were also investigated, since it is necessary, for normal cell turnover and regeneration between lactations. Thus average length variation can affect future production level and, eventually, udder health (Collier *et al.*, 2012). Watters *et al.* (2008) refer that cows submitted to a dry period between 60 and 69 days produced more milk in the subsequent lactation than those given a dry period under 40 days. In the present study it was found that dry periods between 60 and 70 days (which was the maximum dry period length observed in studied herds) increased the probability of isolation of *S. non-agalactiae* ( $p < 0.05$ ). Dry periods shorter than 45 days were less likely than the previous, to increase the probability of isolation. This finding should be cautiously regarded as a protective determinant factor, since the use of this dry period duration reflects the reality of only 9 farms. Furthermore, Church *et al.* (2008) and Pinedo *et al.* (2011) reported that no difference in somatic cell count and cases of mastitis were found for shorted dry periods (from 30 to 34) versus 45-, 55-, and 60-dry periods. In the present study, difference of duration of the dry period influenced the occurrence of mastitis by *S. non-agalactiae*, being the dry period of 45 to 60 days better to prevent infection by this agent.

Possible effects of teat disinfection practices were also investigated, although the composition of the dipping solution was not assessed. It was observed that the simultaneous use of pre- and post-milking teat dipping, was associated with a higher probability of isolating *Streptococcus non-agalactiae* ( $p < 0.05$ ). In contrast, absence of teat disinfection in milking was associated with decreasing isolation probability. Such result was equally found in the analysis of the relationship between isolation of environmental agents and use of teat disinfections ( $p < 0.01$ ). Post-dipping is

used as a control measure of contagious mastitis for several years, whereas pre-dipping, more recent, is used as a control measure in the case of environmental mastitis (Morn, 2009). Teat dipping in solutions composed by, for instance sodium hypochlorite, iodine, chlorhexidine and chorine, decreases the population of microorganisms on the teat skin. At the end of milking, debris of dipping should be discarded and the container should be cleaned before the next milking to avoid contamination. In addition, components of the dipping solution may be inactivated with organic material and thus it is important that the dipping container does not become contaminated with feces (Blowery & Edmondson, 2010). The results obtained in this study, can be explain by bad dipping practices (can act as a source of infection if the product is contaminated at the time of administration when both teat disinfectants are used) or by the fact that dipping practices were being used in the most affected farms.

To assess the influence of the season, the year was divided into four trimesters, starting in January. Significant association ( $p < 0.05$ ) was found between the fourth trimester and increased frequency of *S. uberis* isolates, i.e. between October and December. The first trimester of the year (January-March) followed. The second trimester corresponding to the spring season was the one where fewer *S. uberis* isolates were found.

Similarly to *S. uberis* the frequency of isolates of *Enterococcus* and environmental agents, was significantly higher ( $p < 0.01$ ) in the fourth trimester (from October to December), followed by the first trimester (January until March). The summer season (third trimester) was the period of the year when the frequency of isolates of these agents was lower.

Described results of increased frequency of isolation during the fourth and first trimester can be explained by the recognized effects of temperature and humidity over mastitis (Morse *et al.*, 1988). Indeed, the fourth and first trimester of the year correspond to the months of greater humidity in São Miguel. Although the rain is more or less constant throughout the year in the island, it is more intensive during winter, i.e. from October until March. Similarly, Lopez-Benavides *et al.* (2007) reported the lowest concentration of *S. uberis* in several ecological niches of a dairy farm of New Zealand during the summer season.

Finally, concerning the remaining factors, namely the type of milking machine, the existence of cowshed and the location of farms no significant statistical association was observed. The type of milking machine was used as a variable in the study because a large proportion of producers

use mobile outdoor milking equipment's, given that the cows are at pasture. This is different from the reality observed in the farms of Portugal mainland.

The present study has some limitations, namely due to the impossibility of performing the questionnaires in all of the 133 holdings that provided milk samples to the AJAM laboratory throughout the year of 2013. It was further difficult to engage the producers to answer all of the parameters abridged by the questionnaire. Concerning the laboratory records, the major limitations were the absence of information regarding the patterns of resistance of isolated agents. Comparison with previous national and foreign studies was also hampered by the different ABs employed in the susceptibility testing assays.

The major outcomes of this study were: 1) higher percentage of environmental agents isolated in the analyzed milk samples; 2) isolation of a significant number of contagious agents with a good AB response; 3) association of teat dipping practices and internal teat sealant to a higher percentage of isolation of some agents, contrarily to what might be expected. The findings of the present study suggest that an improvement of the holdings hygiene along with better compliance with good milking practices might contribute to a decrease of mastitis occurrence.

Thus, further work is needed to characterize the features of the Azorean holdings and herds that can sustain the high frequency of occurrence of mastitis caused by environmental agents. It would also be important to determine the resistance patterns and identify the agents involved not only in clinical mastitis, as well as in subclinical mastitis. Other significant aspects that deserve further attention are determination of the risk factors for the occurrence of mastitis, based particularly on the characteristic management of the holdings of São Miguel, hygiene scores, compliance with good milking practices and the influence of different dry periods. By targeting these aspects, better and more effective mastitis control plans could be implemented with the aim to provide a higher quality and valorization of Azorean milk and its derivatives.

## **ACKNOWLEDGMENTS**

The authors are grateful to the *Associação de Jovens Agricultores Micaelenses*, particularly to Dra. L. Soares and Dr. T. Viveiros and laboratory technician R. Sardinha for providing data from the laboratory records and for assisting in the collection of questionnaire data.

The authors also acknowledge Prof. Dr. N. Carolino for the support in the statistical evaluation of the data.

#### **CONFLICT OF INTEREST STATEMENT**

None of the authors of this paper has a financial or personal relationship with other people or organizations that could inappropriately influence or bias the content of the paper.

#### **REFERENCES:**

Aires T., 2010. Mastites em bovinos. Caracterização etiológica, padrões de sensibilidade e implementação de programas de qualidade do leite em explorações do Entre-Douro e Minho, tese de mestrado. Localidade, Universidade Técnica de Lisboa.

Berry E., Hillerton J., 2002. The Effect of an Intramammary Teat Seal on New Intramammary Infections. *Journal of Dairy Science* 85(10), 2512-2520.

Bexiga R., Pereira H., Pereira O., Leitão A., Carneiro C., Ellis K., Vilela C., 2010. Observed reduction in recovery of *Corynebacterium* spp. from bovine milk samples by use of a teat cannula. *Journal of Dairy Science* 78, 9-14.

Blowery R. and Edmondson P. (Eds), 2010. Mastitis Control in Dairy Herds. 2<sup>nd</sup> Ed. CAB International, United Kingdom, pp. 33-59.

Church G., Fox L., Gaskins C., Hancock D., Gay J., 2008. The Effect of a Shortened Dry Period on Intramammary Infections During the Subsequent Lactation. *Journal of Dairy Science* 91(11), 4219-4225.

Collier R., Annen-Dawson E., Pezeshki A., 2012. Effects of continuous lactation and short dry periods on mammary function and animal health. *Animal* 6(3), 403-414.

Guérin-Faubleé V., Tardy F., Bouveron C., Carret G., 2002. Antimicrobial susceptibility of *Streptococcus species* isolated from clinical mastitis in dairy cows. International Journal of antimicrobial Agents 19, 219-226.

Huscley J., Green M., Green L., Bradley A., 2002. Evaluation of the Efficacy of an Internal teat Sealer during the dry period. Journal of dairy Science 85(3), 551-556.

Lopez-Benavides M., Williamson J., Pullinger G., Lacy-Hulbert S., Cursons R., Leigh J., 2007. Field Observations on the Variation of *Streptococcus uberis* Populations in a Pasture-Based Dairy Farm. Journal of Dairy Science 90, 5558-5566.

Melchior M., Osch M., Graat R., Duijkeren E., Mevius D., Nielen M., Gaastra W., Fink-Gremmels J., 2009. Biofilm formation and genotyping of *Staphylococcus aureus* bovine mastitis isolates: Evidence for lack of penicillin-resistance in *Agr*-type II strains. Veterinary Microbiology 137, 83-89.

Merl K., Abdulmawjood A., Lämmle C., Zschöck M., 2003. Determination of epidemiological relationships of *Streptococcus agalactiae* isolated from bovine mastitis. FEMS Microbiology Letters 226, 87-92.

Morn D., 2009. Mammary Gland Health and Disorders. In: Smith B. (Ed.) Large Animal Internal Medicine, 4<sup>th</sup> Ed. Unites States, Mosby Elsevier, pp. 1112-1143.

Morse D., DeLorenzo M., Wilcox C., Collier R., Natzke R., Bray R., 1988. Climatic Effects on Occurrence of Clinical Mastitis. Journal of Dairy Science 71, 848-853.

Peixoto R., França C., Júnior A., Veschi J., Costa M., 2010. Etiologia e perfil de sensibilidade antimicrobiana dos isolados bacterianos da mastite em pequenos ruminantes e concordância de técnicas empregadas no diagnóstico. Pesquisa Vet Brasileira 30(9), 735-740.



Pinedo P., Risco C., Melendez P., 2011. A retrospective study on the association between different lengths of the dry period and subclinical mastitis, milk yield, reproductive performance, and culling in Chilean dairy cows. *Journal of Dairy Science* 94(1), 106-115.

Pinho L., Thompson G., Machado M., Carvalheira J., 2013. Management practices associated with the bulk tank milk prevalence of *Mycoplasma* spp. in dairy herds in Northwestern Portugal. *Preventive Veterinary Medicine* 108, 21-27.

Ponte R., 2009. Estimação da função produção e rendimento de leite por animal aplicação da função gama incompleta à ilha de São Miguel, tese de mestrado. Localidade, Universidade dos Açores.

Pyörälä S., 2009. Treatment of mastitis during lactation. *Irish Veterinary Journal* 62, 40-44.

Rangel P. and Marin J., 2009. Analysis of *Escherichia coli* isolated from bovine mastitic milk. *Pesquisa Vet Brasileira* 29(5), 363-368.

Rato M., Bexiga R., Florindo C., Cavaco L., Vilela C., Santos-Sanches I., 2013. Antimicrobial resistance and molecular epidemiology of streptococci from bovine mastitis. *Veterinary Microbiology* 161, 286-294.

Rocha B., 2013. Antibacterial Resistance Patterns and over time trends of major mastitis pathogens from dairy cattle in Portugal, tese de mestrado. Localidade, Universidade do Porto.

S.R.E.A. (Serviço Regional de Estatística dos Açores), 2014. Entrega de leite na fábrica. Acedido em: 28/03/2014. Disponível em: <http://estatistica.azores.gov.pt/upl/%7B8b054a37-283c-4571-93fd-0f24a5de719e%7D.htm>.

Saini V., McClure J., Léger D., Keefe G., Scholl D., Morch D., Barkema H., 2012. Antimicrobial resistance profiles of common mastitis pathogens on Canadian dairy farms. *Journal of Dairy Science* 95, 4319-4332.

SAS Institute, 2004. SAS 9.1.2 for Microsoft Windows. SAS International, Heidelberg, Germany.  
Seegers H., Fourichon C., Beaudeau F., 2003. Production effects related to mastitis and mastitis economics in dairy cattle herds. *Veterinary Research* 34, 475-491.

Silva E., Siqueira A., Martins J., Ferreira W., Silva N., 2004. Identification and in vitro Antimicrobial susceptibility of *Staphylococcus* species isolated from goat mastitis in the Northeast of Brazil. *Small Ruminant Research* 55, 45-49.

Sousa R., 2007. Clínica de Espécies Pecuárias, tese de mestrado. Localidade, Universidade de Trás os Montes e Alto Douro.

Spanamberg A., Sanches E., Santurio J., Ferreira L., 2009. Mastite micótica em ruminantes causada por leveduras. *Ciência Rural* 39(1), 282-290.

Thorberg B., Danielsson-Tham M., Emanuelson U., Waller K., 2009. Bovine subclinical mastitis caused by different types of coagulase-negative Staphylococci. *Journal of Dairy Science* 92, 4962-4970.

Watters R., Guenther J., Brickner A., Rastani R., Crump P., 2008. Effects of Dry Period Length on Milk Production and Health of Dairy Cattle. *Journal of Dairy Science* 91(7), 2595-2603.

Zadoks R., 2007. Sources and epidemiology of *Streptococcus uberis*, with special emphasis on mastitis in dairy cattle. *Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* 2(30), 15.

Zadoks R., Allore H., Barkema H., Sampimon O., Grohn Y., Schukken H., 2001. Analysis of an Outbreak of *Streptococcus uberis* Mastitis. Journal of Dairy Science 84, 590-599.

# ANNEX

# Annex 1

**Guide for authors of the journal *Research in Veterinary Science*** (retrieved from: <http://www.elsevier.com/journals/research-in-veterinary-science/0034-5288/guide-for-authors>):

- *Research in Veterinary Science* publishes original contributions and review articles on research concerning the health and disease of animals, including studies in comparative medicine.

#### **Types of contribution**

1. Original research papers (Regular Papers)
2. Short Communications
3. Review articles

**Original research papers** should report the results of original research. The material should not have been previously published elsewhere, except in a preliminary form.

**Short Communications** should not exceed 1600 words and include no more than two tables or figures. They should have an abstract but no other divisions. Typescripts should be clearly marked Short Communication.

**Review articles** Review articles on veterinary topics are invited for publication. They should give an update on recent advances in a particular field and be targeted at research veterinarians who are not necessarily working in the same field. The length should not exceed 4000 words.

#### **Submission of manuscripts**

Submission to *Research in Veterinary Science* now proceeds online via Elsevier Editorial System -<http://ees.elsevier.com/rvsc>. Authors will be guided step-by-step through uploading files directly from their computers. Authors should select a set of classifications for their papers from a given list, as well as a category designation (Original Research Paper, Short Communication, and so on). Electronic PDF proofs will be automatically generated from uploaded files, and used for subsequent reviewing.

Authors should send queries concerning the submission process or journal procedures to [AuthorSupport@elsevier.com](mailto:AuthorSupport@elsevier.com). Authors can check the status of their manuscript within the review procedure using Elsevier Editorial System.

Authors submitting hard copy papers will be asked to resubmit using Elsevier Editorial System.

Submission of an article is understood to imply that the article is original and is not being considered for publication elsewhere. Submission also implies that all authors have approved the paper for release and are in agreement with its content. Upon acceptance of the article by the journal, the author(s) will be asked to transfer the copyright of the article to the Publisher. This transfer will ensure the widest possible dissemination of information.

All authors should have made substantial contributions to all of the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, (3) final approval of the version to be submitted.

### **Acknowledgements**

All contributors who do not meet the criteria for authorship as defined above should be listed in an acknowledgements section. Examples of those who might be acknowledged include a person who provided purely technical help, writing assistance, or a department chair who provided only general support. Authors should disclose whether they had any writing assistance and identify the entity that paid for this assistance.

### **Conflict of interest**

At the end of the text, under a subheading "Conflict of interest statement" all authors must disclose any financial and personal relationships with other people or organisations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding.

### **Role of the funding source**

All sources of funding should be declared as an acknowledgement at the end of the text. Authors should declare the role of study sponsors, if any, in the study design, in the collection, analysis and interpretation of data; in the writing of the manuscript; and in the decision to submit the manuscript for publication. If the study sponsors had no such involvement, the authors should so state.

**Open Access** This journal offers authors two choices to publish their research;

*1. Open Access*

- Articles are freely available to both subscribers and the wider public with permitted reuse
- An Open Access publication fee is payable by authors or their research funder

*2. Subscription*

- Articles are made available to subscribers as well as developing countries and patient groups through our access programs (<http://www.elsevier.com/access>)
- No Open Access publication fee

All articles published Open Access will be immediately and permanently free for everyone to read and download. Permitted reuse is defined by your choice of one of the following Creative Commons user licenses:

*Creative Commons Attribution-Non Commercial-ShareAlike (CC BY-NC-SA)*: for non-commercial purposes, lets others distribute and copy the article, to create extracts, abstracts and other revised versions, adaptations or derivative works of or from an article (such as a translation), to include in a collective work (such as an anthology), to text and data mine the article, as long as they credit the author(s), do not represent the author as endorsing their adaptation of the article, do not modify the article in such a way as to damage the author's honor or reputation, and license their new adaptations or creations under identical terms (CC BY NC SA).

*Creative Commons Attribution-NonCommercial-NoDerivs (CC-BY-NC-ND)*: for non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

*Creative Commons Attribution (CC-BY)*: available only for authors funded by organizations with which we have established an agreement with. For a full list please see <http://www.elsevier.com/journal-authors/funding-body-agreements>.



To provide Open Access, this journal has a publication fee which needs to be met by the authors or their research funders for each article published Open Access. Your publication choice will have no effect on the peer review process or acceptance of submitted articles.

The Open Access publication fee for this journal is \$USD 2,500, excluding taxes.

Learn more about Elsevier's pricing policy [www.elsevier.com/openaccesspricing](http://www.elsevier.com/openaccesspricing).

**Language Editing:** [Elsevier's Authors Home](#) provides details of some companies who can provide English language and copyediting services to authors who need assistance *before* they submit their article or *before* it is accepted for publication. Authors should contact these services directly. For more information about language editing services, please email [authorsupport@elsevier.com](mailto:authorsupport@elsevier.com).

## **Ethics**

Before papers describing animal studies are accepted for publication in *Research in Veterinary Science*, the authors must satisfy the editors that the work conformed to appropriate ethical standards. Whether or not a particular piece of work is accepted for publication will be decided by the editors whose decision will be final.

The authors should provide written assurances that: (i) The project underwent ethical review and was given approval by an institutional animal care and use committee or by appropriately qualified scientific and lay colleagues. (ii) The care and use of experimental animals complied with local animal welfare laws, guidelines and policies.

The editors expect authors to have adhered to the following general principles: (i) Alternative procedures that replace the use of animals should be used if possible. Where this is not possible, the animals used should be carefully selected to be the least sentient species possible and of an appropriate strain. (ii) The minimum number of animals should be used consistent with achieving the scientific objectives of the study. (iii) Pain and distress should be minimised by the use of humane endpoints, sedation, anaesthesia, analgesia and post-operative care. (iv) Access to veterinary care must be available at all times. (v) Investigators and personnel that

care for and use animals must be trained and possess relevant expertise and training that should be updated regularly. (vi) If animals have to be killed, this should be done humanely according to local euthanasia regulations, such as the Home Office guidelines in the UK or guidelines of the American Veterinary Association Panel on Euthanasia.

### **Title**

Papers should be headed with the full title, the initials and surnames of the authors, and the name and address of the institution where the work was carried out. The full telephone number, Fax number and e-mail address of the corresponding author should also be provided.

### **Form of Papers**

a) Abstract (not more than 150 words), self-contained and embodying the main conclusions. It should note the relevance to veterinary science as well as the aims and objectives of the work. Sentences such as 'the results are discussed', which merely describe the paper, are not allowed.

b) Keywords. Please supply a list of up to six keywords that describe the paper.

c) Introduction.

d) Materials and methods employed.

e) Results, as concise as possible. Text, tables and figures illustrating the same data will rarely be permitted.

f) Discussion and conclusions.

g) Acknowledgements.

h) References.

i) Manuscripts should have **numbered lines**, with wide margins and **double spacing**, throughout, i.e. also for abstracts, footnotes and references. **Every page of the manuscripts, including the title page, references, tables, etc., should be numbered.** However, in the text no reference should be made to page numbers; if necessary one may refer to sections. Avoid excessive usage of italics to emphasize part of the text.

**Abbreviation and symbols:** Authors are asked to explain each scientific abbreviation at its first occurrence in their papers; for example, complement fixation test (CFT). The policy of the journal with respect to units and symbols is that SI (System International) symbols should be used.

## **References**

1. All publications cited in the text should be presented in a list of references following the text of the manuscript. The manuscript should be carefully checked to ensure that the spelling of author's names and dates are exactly the same in the text as in the reference list.
2. In the text refer to the author's name (without initial) and year of publication, followed - if necessary - by a short reference to appropriate pages. Examples: "Since Peterson (1988) has shown that..." "This is in agreement with results obtained later (Kramer, 1989, pp. 12-16)".
3. If reference is made in the text to a publication written by more than two authors the name of the first author should be used followed by "et al." This indication, however, should never be used in the list of references. In this list names of first author and co-authors should be mentioned.
4. References cited together in the text should be arranged chronologically. The list of references should be arranged alphabetically on author's names, and chronologically per author. If an author's name in the list is also mentioned with co-authors the following order should be used: publications of the single author, arranged according to publication dates - publications of the same author with one co-author - publications of the author with more than one co-author. Publications by the same author(s) in the same year should be listed as 1974a, 1974b, etc.
5. Use the following system for arranging your references:
  - a. For periodicals

Minamoto, T., Honjo, M.N., Yamanaka, H., Tanaka, N., Itayama, T., Kawabata, Z., 2011. Detection of cyprinid herpesvirus-3 DNA in lake plankton. *Research in Veterinary Science* 90, 530-532.

Castillo, V.A., Gomez, N.V., Lalia, J.C., Cabrera Blatter, M.F., Garcia, J.D., 2008a. Cushing's disease in dogs: Cabergoline treatment. *Research in Veterinary Science* 85, 26-34.

b. For books

Blaha, T. (Ed.), 1989. *Applied Veterinary Epidemiology*. Elsevier, Amsterdam, 344 pp.

c. For multi-author books

Wilson, M.B., Nakane, P.K., 1978. Recent developments in the periodate method of conjugating horseradish peroxidase (HRPO) to antibodies. In: Knapp, W., Holubar, K., Wick, G. (Eds.), *Immunofluorescence and Related Staining Techniques*. North Holland, Amsterdam, pp. 215-2246.

6. Please do not abbreviate the journal title names e.g. *Research in Veterinary Science* and not *Res Vet Sci*.

7. In the case of publications in any language other than English, the original title is to be retained. However, the titles of publications in non-Latin alphabets should be transliterated, and a notation such as "(in Russian)" or "(in Greek, with English abstract)" should be added.

8. Work accepted for publication but not yet published should be referred to as "in press".

9. References concerning unpublished data and "personal communications" should not be cited in the reference list but may be mentioned in the text.

10. Web references may be given. As a minimum, the full URL is necessary. Any further information, such as Author names, dates, reference to a source publication and so on, should also be given.

11. Articles available online but without volume and page numbers may be referred to by means of their Digital Object identifier (DOI) code.

## **Illustrations**

1. All illustrations (line drawings and photographs) should be submitted as separate files, preferably in TIFF or EPS format.

2. Illustrations should be numbered according to their sequence in the text. References should be made in the text to each illustration.

3. Illustrations should be designed with the format of the page of the journal in mind. Illustrations should be of such a size as to allow a reduction of 50%.

4. Lettering should be big enough to allow a reduction of 50% without becoming illegible. Any lettering should be in English. Use the same kind of lettering throughout and follow the style of the journal.

5. If a scale should be given, use bar scales on all illustrations instead of numerical scales that must be changed with reduction.

6. Each illustration should have a caption. The captions to all illustrations should be typed on a separate sheet of the manuscript.

7. Explanations should be given in the figure legend(s). Drawn text in the illustrations should be kept to a minimum.

8. Photographs are only acceptable if they have good contrast and intensity.

9. If you submit usable colour figures, Elsevier would ensure that these figures appeared free-of-charge in colour in the electronic version of your accepted paper, regardless of whether or not these illustrations are reproduced in colour in the printed version. Colour illustrations can only be included in print if the additional cost of reproduction is contributed by the author: you would receive information regarding the costs from Elsevier after receipt of your accepted article.

Please note that because of technical complications which may arise by converting colour figures to 'grey scale' (for the printed version, should you not opt for colour in print), you should submit in addition usable black and white figures corresponding to all colour illustrations.

10. Advice on the preparation of illustrations can be found at the following URL: <http://www.elsevier.com/artworkinstructions>

### **Preparation of supplementary data**

Elsevier now accepts electronic supplementary material to support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, movies, animation sequences, high-resolution images, background datasets,

sound clips and more. Supplementary files supplied will be published online alongside the electronic version of your article in Elsevier web products, including

ScienceDirect: <http://www.sciencedirect.com>. In order to ensure that your submitted material is directly usable, please ensure that data is provided in one of our recommended file formats.

Authors should submit the material together with the article and supply a concise and descriptive caption for each file.

### **Tables**

1. Authors should take notice of the limitations set by the size and lay-out of the journal. Large tables should be avoided. Reversing columns and rows will often reduce the dimensions of a table.
2. If many data are to be presented, an attempt should be made to divide them over two or more tables.
3. Tables should be numbered according to their sequence in the text. The text should include references to all tables.
4. Each table should occupy a separate page of the manuscript. Tables should never be included in the text.
5. Each table should have a brief and self-explanatory title.
6. Column headings should be brief, but sufficiently explanatory. Standard abbreviations of units of measurement should be added between parentheses.
7. Vertical lines should not be used to separate columns. Leave some extra space between the columns instead.
8. Any explanation essential to the understanding of the table should be given as a footnote at the bottom of the table.

### **Copyright**

If excerpts from other copyrighted works are included, the Author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by Authors in these cases: contact Elsevier's Rights Department,

Oxford, UK: phone (+1) 215 239 3804 or +44(0)1865 843830, fax +44(0)1865 853333, e-mail [healthpermissions@elsevier.com](mailto:healthpermissions@elsevier.com). Requests may also be completed online via the Elsevier homepage <http://www.elsevier.com/permissions>.

Material in unpublished letters and manuscripts is also protected and must not be published unless permission has been obtained.

### **Authors Rights**

As an author you (or your employer or institution) may do the following:

- make copies (print or electronic) of the article for your own personal use, including for your own classroom teaching use
- make copies and distribute such copies (including through e-mail) of the article to research colleagues, for the personal use by such colleagues (but not commercially or systematically, e.g., via an e-mail list or list server)
- post a pre-print version of the article on Internet websites including electronic pre-print servers, and to retain indefinitely such version on such servers or sites
- post a revised personal version of the final text of the article (to reflect changes made in the peer review and editing process) on your personal or institutional website or server, with a link to the journal homepage (on [elsevier.com](http://elsevier.com))
- present the article at a meeting or conference and to distribute copies of the article to the delegates attending such a meeting
- for your employer, if the article is a 'work for hire', made within the scope of your employment, your employer may use all or part of the information in the article for other intra-company use (e.g., training)
- retain patent and trademark rights and rights to any processes or procedure described in the article
- include the article in full or in part in a thesis or dissertation (provided that this is not to be published commercially)

- use the article or any part thereof in a printed compilation of your works, such as collected writings or lecture notes (subsequent to publication of your article in the journal)
- prepare other derivative works, to extend the article into book-length form, or to otherwise re-use portions or excerpts in other works, with full acknowledgement of its original publication in the journal

### **Funding body agreements and policies**

Elsevier has established agreements and developed policies to allow authors who publish in Elsevier journals to comply with potential manuscript archiving requirements as specified as conditions of their grant awards. To learn more about existing agreements and policies please visit <http://www.elsevier.com/fundingbodies>).

### **Proofs**

One set of page proofs in PDF format will be sent by e-mail to the corresponding author (if we do not have an e-mail address then paper proofs will be sent by post). Elsevier now sends PDF proofs which can be annotated; for this you will need to download Adobe Reader version 7 available free from <http://www.adobe.com/products/acrobat/readstep2.html>. Instructions on how to annotate PDF files will accompany the proofs. The exact system requirements are given at the Adobe site: <http://www.adobe.com/products/acrobat/acrrsystemreqs.html#70win>. If you do not wish to use the PDF annotations function, you may list the corrections (including replies to the Query Form) and return to Elsevier in an e-mail. Please list your corrections quoting line number. If, for any reason, this is not possible, then mark the corrections and any other comments (including replies to the Query Form) on a printout of your proof and return by fax, or scan the pages and e-mail, or by post. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. We will do everything possible to get your article published quickly and accurately. Therefore, it is important to ensure that all of your corrections are sent back to us in one communication: please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

### **Author Enquiries**



For enquiries relating to the submission of articles (including electronic submission where available) please visit the journal's homepage at <http://www.elsevier.com/locate/rvsc>. This also provides the facility to track accepted articles and set up e-mail alerts to inform you of when an article's status has changed.

Contact details for questions arising after acceptance of an article, especially those relating to proofs, are provided after registration of an article for publication.

### **Offprints**

The corresponding author will, at no cost, be provided with a PDF file of the article via e-mail. The PDF file is a watermarked version of the published article and includes a cover sheet with the journal cover image and a disclaimer outlining the terms and conditions of use.

### ***Research in Veterinary Science has no page charges***

### **AudioSlides**

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. More information and examples are available at <http://www.elsevier.com/audioslides>. Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

# Annex 2

**Estudo retrospectivo sobre agentes de mastites e a sua  
sensibilidade a antimicrobianos em explorações de bovinos de  
leite de S. Miguel (Açores)**

# **Questionário**

## **NOTAS PRÉVIAS**

Os dados recolhidos por este questionário destinam-se exclusivamente a fins académicos, especificamente no âmbito da Tese de Mestrado em Medicina Veterinária da aluna da Escola Universitária Vasco da Gama Andreia Filipa Gomes Vieira, sendo que a confidencialidade dos dados recolhidos é assegurada.

Os dados são recolhidos após o consentimento expresso dos respectivos proprietários, previamente informados sobre a natureza e objectivos do estudo.

### IDENTIFICAÇÃO DA EXPLORAÇÃO

- A. Nome do proprietário: \_\_\_\_\_
- B. Local da exploração (Concelho): \_\_\_\_\_
- C. Número de animais na exploração: \_\_\_\_\_
- D. Número de vacas em ordenha: \_\_\_\_\_

### CARACTERÍSTICAS DA EXPLORAÇÃO

- E. Número de vacas com mais que um parto: \_\_\_\_\_
- F. Tipo de exploração:
- Vacas sempre em pastoreio ☐
  - Vacas secas em pastoreio e vacas em lactação estabuladas ☐
  - Vacas sempre estabuladas ☐
  - Estabulação no Inverno ☐
- G. Realiza a recria:
- Sim ☐ Não ☐
- H. Exploração:
- Aberta ☐ Fechada ☐

H.1. Se **ABERTA** adquiriu animais:

- Nos últimos 6 meses ☐
- No último ano ☐
- Há mais de um ano ☐

### MANEIO DAS VACAS SECAS

**I. Duração em média do período de seca:**

> que 60 dias e < que 70 dias ☐

> 70 dias ☐

> que 45 e < que 60 dias ☐

< 45 dias ☐

**J. Realiza a separação de vacas secas das vacas em lactação:**

Sim ☐

Não ☐

**K. Utiliza antibióticos intraamamários à secagem:**

Sempre ☐

Frequentemente ☐

Raramente ☐

Só em caso de mamites identificadas ☐

Nunca ☐

**L. Utiliza selantes internos a secagem:**

Sempre ☐

Frequentemente ☐

Raramente ☐

Nunca ☐

**M. Realiza a testagem de animais a secagem:**

Sempre ☐

Frequentemente ☐

Raramente ☐

Nunca ☐

12.1 Se **REALIZA** testagem, qual:

Isolamento e TSA ☐

TCM ☐

**N. Local do parto:**

Maternidade individual ☐

Parque das secas ☐

Parque das vacas em lactação ☐

Outro: \_\_\_\_\_

**HIGIENE NA ORDENHA**

**O. Utiliza luvas durante a ordenha:**

Sim ☐ Não ☐

**P. Utiliza:**

Pré-dipping ☐

Pós-dipping ☐

Ambos ☐

Nenhum ☐

**Q. Realiza a limpeza do úbere com toalhetes individuais secos:**

Sim ☐ Não ☐

**R. Desinfecção de tetinas entre ordenhas:**

Sim ☐ Não ☐

**S. Ordenha as vacas com problemas de mastites no fim da ordenha:**

Sim ☐ Não ☐

**T. Tipo de ordenha:**

Móvel ☐ Fixa ☐

**U. Nº de pontos de ordenha: \_\_\_\_\_**

**MANEIO ALIMENTAR VACAS SECAS, PRODUÇÃO, NOVILHAS**

**V. Utiliza correctores vitamínicos e minerais nas vacas em lactação:**

Sim ☐ Não ☐

**W. E nas vacas secas:**

Sim ☐

Não ☐